



15-2.6-9.1 NPK PLUS

DESCRIPTION: A PROFESSIONAL HOMOGENEOUS 10 TO 12 MONTH \diamond CONTROLLED RELEASE NURSERY FERTILIZER FOR WOODY ORNAMENTALS, PROPAGATION & FOLIAGE PLANTS. ALL OF THE PRIMARY NUTRIENTS, MAGNESIUM, SULFUR, AND MICRONUTRIENTS IN **POLYON NPK PLUS MICROS 15-2.6-9.1** ARE COMBINED WITHIN EACH UNIFORM COATED PELLET, INSURING PRECISE DISTRIBUTION AND RELEASE.

BENEFITS:

- 15-2.6-9.1 NPK PLUS MICROS provides the improved safety of POLYON® Reactive Layers Coating (RLC) controlled release technology.
- Release of nutrients with POLYON® is predictable and reliable. The coating has been precisely applied to ensure the safety and effectiveness of each granule.
- Release of nutrients is not significantly affected by media type, moisture level, pH, or modified activity.



SOIL/MEDIA TEMPERATURE RELEASE RATES

50°F 10.0°C = 14-15 months
60°F 15.5°C = 12-13 months
70°F 21.0°C = 10-11 months \diamond
80°F 26.5°C = 8-9 months

POLYON 15-6-12 NPK PLUS GUARANTEED ANALYSIS:

U.S. STANDARD

TOTAL NITROGEN (N)*	15.00%
8.30% Ammoniacal Nitrogen	
6.70% Nitrate Nitrogen	
AVAILABLE PHOSPHATE (P ₂ O ₅)*	6.00%
SOLUBLE POTASH (K ₂ O)*	12.00%
Magnesium (Mg)*	1.45%
Sulfur (S)*	6.00%
Copper (Cu)*	0.05%
Iron (Fe)*	0.40%
0.40% Chelated Iron	
Manganese (Mn)*	0.10%
Molybdenum (Mo)*	0.0199%
Zinc (Zn)*	0.05%

Derived from Polymer-Coated Ammonium Nitrate, Polymer-Coated Ammonium Phosphate, Polymer-Coated Sulfate of Potash, Polymer-Coated Magnesium Sulfate, Polymer-Coated Copper Sulfate, Polymer-Coated Iron EDTA, Polymer-Coated Manganese Sulfate, Polymer-Coated Sodium Molybdate, and Polymer-Coated Zinc Sulfate.

POLYON® is a registered trademark owned by Agrium and used under license.

*All nutrients have been polymer-coated to provide 15.00% coated slow release nitrogen (N), 6.00% coated slow release available phosphate (P₂O₅), 12.00% coated slow release soluble potash (K₂O), 1.45% coated slow release magnesium (Mg), 6.00% coated slow release sulfur (S), 0.05% coated slow release copper (Cu), 0.40% coated slow release iron (Fe), 0.10% coated slow release manganese (Mn), 0.0199% coated slow release molybdenum (Mo), and 0.05% coated slow release zinc (Zn).

POLYON 15-2.6-9.1 NPK PLUS GUARANTEED ANALYSIS:

ELEMENTAL

TOTAL NITROGEN (N)**	15.00%
8.30% Ammoniacal Nitrogen	
6.70% Nitrate Nitrogen	
TOTAL PHOSPHORUS (P)**	2.60%
TOTAL POTASSIUM (K)**	9.90%
Magnesium (Mg)**	1.45%
Sulfur (S)**	6.00%
Copper (Cu)**	0.05%
Iron (Fe)**	0.40%
0.40% Chelated Iron	
Manganese (Mn)**	0.10%
Molybdenum (Mo)**	0.0199%
Zinc (Zn)**	0.05%

Derived from Polymer-Coated Ammonium Nitrate, Polymer-Coated Ammonium Phosphate, Polymer-Coated Sulfate of Potash, Polymer-Coated Magnesium Sulfate, Polymer-Coated Copper Sulfate, Polymer-Coated Iron EDTA, Polymer-Coated Manganese Sulfate, Polymer-Coated Sodium Molybdate, and Polymer-Coated Zinc Sulfate.

POLYON® is a registered trademark owned by Agrium and used under license.

**All nutrients have been polymer-coated to provide 15.00% coated slow release nitrogen (N), 2.60% coated slow release total phosphorus (P), 9.90% coated slow release total potassium (K), 1.45% coated slow release magnesium (Mg), 6.00% coated slow release sulfur (S), 0.05% coated slow release copper (Cu), 0.40% coated slow release iron (Fe), 0.10% coated slow release manganese (Mn), 0.0199% coated slow release molybdenum (Mo), and 0.05% coated slow release zinc (Zn).

APPLICATION RATES: (Call for rates on larger containers.)

Use **LOW** rate for low feeding, sensitive plants or under high soil temperatures.

Use **MEDIUM** rate for medium to moderately heavy feeding plants.

Use **HIGH** rate only for heavy feeding hardy plants.

These application rates are based on the average temperature at the fertilizer location of 70° F (21.0°C).

Increase fertilizer application rates by 20% if average monthly temperatures are lower than 60°F (15.5°C).

Lower application rates by 20% if average monthly temperatures are greater than 80°F (26.5°C).

CONVERSION TABLE	DRY MEASURE		
	Level Measure	Grams	Oz.(Wt.)
	1 teaspoon (tsp.)	5.7	0.20
	1 tablespoon (tblsp.)	17.8	0.63
	1/4 cup	56.1	1.98
	1/2 cup	121.3	4.28

CONVERSION TABLE	POLYON SPOONS					
	Size	Grams	Oz. (Wt.)	Size	Grams	Oz.(Wt.)
	1	10.5	0.37	5	37.8	1.33
	2	15.6	0.55	6	54.2	1.91
	3	21.1	0.74	7	71.3	2.52
	4	26.4	0.93	8	86.0	3.03

TOPDRESS CONTAINER: Plant Nutrient Requirements / Uniformly apply (topdress) product onto the container surface using the amounts listed below.

VOLUME (gal.)	DIAMETER	LOW	MEDIUM	HIGH	DIAMETER (mm)	LOW	MEDIUM	HIGH
1 gallon	6 inches	10 g	15 g	20 g	100mm	2.0 g	3.0 g	4.0 g
2 gallons	8 inches	20 g	30 g	40 g	125mm	4.0 g	6.0 g	8.0 g
3 gallons	10 inches	40 g	60 g	80 g	150mm	7.0 g	10.0 g	13.0 g
5 gallons	12 inches	60 g	90 g	120 g	175mm	12.0 g	18.0 g	24.0 g
7 gallons	14 inches	85 g	130 g	175 g	200mm	18.0 g	27.0 g	36.0 g
10 gallons	17 inches	150 g	225 g	300 g	250mm	40.0 g	60.0 g	80.0 g
15 gallons	18 inches	200 g	300 g	400 g	300mm	55.0 g	80.0 g	105.0 g

INCORPORATION: Plant Nutrient Requirements / Uniformly mix (incorporate) nursery fertilizer into potting media as follows:

POUNDS PER CUBIC YARD	LOW 10	MED 15	HIGH 20	KILOGRAMS PER CUBIC METRES	LOW 6	MED 9	HIGH 12
-----------------------	--------	--------	---------	----------------------------	-------	-------	---------

PLANTING BED: FIELD / Plant Nutrient Requirements (incorporate if possible or use lower rates) as follows:

POUNDS PER 100 SQ.FT.	LOW 4	MED 7	HIGH 10	KILOGRAMS PER 100 SQ. METRES	LOW 20	MED 32	HIGH 48
-----------------------	-------	-------	---------	------------------------------	--------	--------	---------

Item# 779584_15612npkmax_3-5-09

APPLICATION PRECAUTIONS:

- Trial before use of this product under your local growing conditions, application methods, and desired rates. Avoid application to plants under stress.
- If mixed media is not used within 1 week, leach thoroughly before using.
- Product left in media for more than 1 week will lose longevity resulting in reduced release time and wasted controlled release fertilizer.
- Avoid the use of media processing equipment that could change the integrity of RLC.
- Avoid mounding of fertilizer against base of plant.
- Iron and other plant nutrients can cause staining of cement.
- Keep away from pools, ponds, and other bodies of water.
- When using potting media with higher cation exchange capacities use lower recommended rates of this formulation.
- When using supplemental liquid feed reduce the rate of this formulation accordingly.
- Do not incorporate into media prior to steam sterilization.
- This product is not recommended for dibble applications.
- To avoid buildup of soluble salts, occasional leaching may be necessary.
- CAUTION: Application of fertilizer materials containing Molybdenum (Mo) may result in forage crops containing levels of Molybdenum (Mo) that are toxic to ruminant animals.